

THE MEAN CURVATURE FLOW APPROACH TO THE SYMPLECTIC ISOTOPY PROBLEM

XIAOLI HAN, JIAYU LI

ABSTRACT. Let M be a Kähler-Einstein surface with positive scalar curvature. If the initial surface is sufficiently close to a holomorphic curve, we show that the mean curvature flow has a global solution and it converges to a holomorphic curve.

1. INTRODUCTION

It was conjectured by Tian [T] that every embedded orientable closed symplectic surface in a compact Kähler-Einstein surface is isotopic to a symplectic minimal surface in a suitable sense. When Kähler-Einstein surfaces are of positive scalar curvature, this was proved for lower degrees by using pseudo-holomorphic curves (cf. [ST], [Sh]). Since 1998, we have been trying to study Tian's conjecture by using the mean curvature flow (c.f. [CT], [CL1], [CL2]). The mean curvature flow was introduced and intensively studied by Huisken ([H], [EH]).

The progress has been made during these years.

Suppose that M is a Kähler-Einstein surface. Let ω be the symplectic form on M and let J be a complex structure compatible with ω . The Riemannian metric $\langle, \rangle$ on M is defined by

$$\langle U, V \rangle = \omega(U, JV).$$

For a compact oriented real surface Σ which is smoothly immersed in M , one defines, following [CW], the kähler angle α of Σ in M by

$$\omega|_{\Sigma} = \cos \alpha d\mu_{\Sigma}$$

where $d\mu_{\Sigma}$ is the area element of Σ in the induced metric from $\langle, \rangle$. We say that Σ is a holomorphic curve if $\cos \alpha \equiv 1$, Σ is a Lagrangian surface if $\cos \alpha \equiv 0$ and Σ is a symplectic surface if $\cos \alpha > 0$.

The evolution equation for the Kähler angle was obtained in [CT] and [CL1] (also see [Wa]). Let J_{Σ_t} be an almost complex structure in a tubular neighborhood of Σ_t on M with

$$\begin{cases} J_{\Sigma_t} e_1 &= e_2 \\ J_{\Sigma_t} e_2 &= -e_1 \\ J_{\Sigma_t} v_1 &= v_2 \\ J_{\Sigma_t} v_2 &= -v_1. \end{cases} \quad (1.1)$$

Key words and phrases. Mean curvature flow, symplectic surface, holomorphic curve.
The research was supported by NSFC .

It is not difficult to check (c.f. [CL1]), with $\bar{\nabla}$ being the covariant derivative of the metric $\langle, \rangle$ on M , that

$$|\bar{\nabla} J_{\Sigma_t}|^2 = |h_{11}^2 + h_{12}^1|^2 + |h_{21}^2 + h_{22}^1|^2 + |h_{12}^2 - h_{11}^1|^2 + |h_{22}^2 - h_{21}^1|^2 \geq \frac{1}{2}|H|^2. \quad (1.2)$$

Proposition 1.1. ([CL1], [CT]) *Suppose that M is a Kähler-Einstein surface with scalar curvature R . Let α be the Kähler angle of the surface Σ_t which evolves by the mean curvature flow. Then*

$$\left(\frac{\partial}{\partial t} - \Delta\right) \cos \alpha = |\bar{\nabla} J_{\Sigma_t}|^2 \cos \alpha + R \sin^2 \alpha \cos \alpha \quad (1.3)$$

where R is the scalar curvature of M .

One corollary of the evolution equation is that, if the initial surface is symplectic, along the mean curvature flow, at every time t the surface is still symplectic.

It is proved in [CLT] that, if the scalar curvature of the Kähler-Einstein surface is positive and the mean curvature flow with initial surface symplectic exists globally, it sub-converges to a holomorphic curve at infinity outside a finite set of points.

To study the global existence of the mean curvature flow with initial surface symplectic, we ([CL1], [CL2]) derived a new monotonicity inequality and proved that the tangent cone at the first blow-up time is flat which implies that there is no Type I singularity (also see [Wa]).

In this paper we will show that, *if the scalar curvature of the Kähler-Einstein surface is positive and the initial surface is sufficiently close to a holomorphic curve, the mean curvature flow has a global solution and it converges to a holomorphic curve.*

The main new point of this paper is the integral estimate of the Kähler angle

$$\int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq \int_{\Sigma_0} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_0 \exp^{-Rt},$$

which implies the estimation of the mean curvature vector on the time space,

$$\int_0^T \int_{\Sigma_t} |H| d\mu_t dt \leq \left(\int_{\Sigma_0} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_0 \right)^{1/2} \frac{\text{Area}(\Sigma_0)^{1/2}}{1 - \exp^{-\frac{R}{2}}}.$$

Using the last inequality, we show that the parabolic density function at any time t along the mean curvature can be dominated by the one at beginning, then our result follows from White's [W] regularity theorem.

The result in this paper gives more evidence that the program should be realized in general.

2. THE MAIN RESULT AND ITS PROOF

We first derive an integral estimation of the Kähler angle along the flow, which yields an L^2 -estimation of the mean curvature vector on the time space.

Proposition 2.1. *Suppose that M is a Kähler-Einstein surface with scalar curvature R . Let α be the Kähler angle of the surface Σ_t which evolves by the mean curvature flow. Suppose that $\cos \alpha(\cdot, 0)$ has a positive lower bound. Then*

$$\int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq C_0 \exp^{-Rt}$$

and

$$\int_t^{t+1} \int_{\Sigma_t} |H|^2 d\mu_t dt \leq C_0 \exp^{-Rt}$$

where C_0 is a constant which depends only on the initial surface, $C_0 = \int_{\Sigma_0} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_0$.

Proof: By applying the parabolic maximum principle to the evolution equation (1.3), one concludes that $\cos \alpha$ remains positive as long as the mean curvature flow has a smooth solution, no matter R is positive, 0 or negative (c.f. [CT], [CL1], [Wa1]). Moreover, we can obtain that

$$\left(\frac{\partial}{\partial t} - \Delta\right) \frac{1}{\cos \alpha} = -\frac{1}{\cos^2 \alpha} (|\bar{\nabla} J_{\Sigma_t}|^2 \cos \alpha + R \sin^2 \alpha \cos \alpha) - \frac{2|\nabla \cos \alpha|^2}{\cos^3 \alpha}.$$

Because ω is closed, we can see that

$$\int_{\Sigma_t} \cos \alpha d\mu_t = \int_{\Sigma_t} \omega$$

is constant under the continuous deformation in t . In fact, for any $t_1 < t_2$, let $\Omega_{t_1, t_2} = \cup_{t_1 \leq t \leq t_2} \Sigma_t$ be the body spread by Σ_t from t_1 to t_2 , since $\partial \Sigma_{t_1} = \partial \Sigma_{t_2} = \emptyset$, by Stokes theorem, one has

$$\int_{\Sigma_{t_2}} \omega - \int_{\Sigma_{t_1}} \omega = \int_{\Omega_{t_1, t_2}} d\omega = 0.$$

We therefore have

$$\begin{aligned} \frac{\partial}{\partial t} \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t &= \frac{\partial}{\partial t} \int_{\Sigma_t} \frac{1}{\cos \alpha} d\mu_t = \int_{\Sigma_t} \left(\frac{\partial}{\partial t} - \Delta\right) \frac{1}{\cos \alpha} d\mu_t - \int_{\Sigma_t} \frac{|H|^2}{\cos \alpha} d\mu_t \quad (2.1) \\ &= - \int_{\Sigma_t} \frac{|\bar{\nabla} J|^2}{\cos \alpha} d\mu_t - \int_{\Sigma_t} R \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t - \int_{\Sigma_t} \frac{|\nabla \cos \alpha|^2}{\cos^3 \alpha} d\mu_t - \int_{\Sigma_t} \frac{|H|^2}{\cos \alpha} d\mu_t. \end{aligned}$$

So,

$$\frac{\partial}{\partial t} \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq -R \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t.$$

It is easy to get that

$$\int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq \int_{\Sigma_0} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_0 \exp^{-Rt}$$

i.e.,

$$\int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq C_0 \exp^{-Rt}.$$

From (2.1) we know that

$$\int_{\Sigma_t} |H|^2 d\mu_t \leq -\frac{\partial}{\partial t} \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t.$$

Integrating the above inequality from t to $t+1$ we obtain that

$$\begin{aligned} \int_t^{t+1} \int_{\Sigma_t} |H|^2 d\mu_t dt &\leq \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t - \int_{\Sigma_{t+1}} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \\ &\leq \int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq C_0 \exp^{-Rt}. \end{aligned}$$

This proves the proposition. Q.E.D.

We derive an L^1 -estimation of the mean curvature vector on the time space.

Proposition 2.2. *Suppose that M is a Kähler-Einstein surface with positive scalar curvature R . Let α be the Kähler angle of the surface Σ_t which evolves by the mean curvature flow. Suppose that $\cos \alpha(\cdot, 0)$ has a positive lower bound. Then*

$$\int_0^T \int_{\Sigma_t} |H| d\mu_t dt \leq (C_0)^{1/2} \frac{\text{Area}(\Sigma_0)^{1/2}}{1 - \exp^{-\frac{R}{2}}}.$$

where the constant C_0 is defined in Proposition 2.1.

Proof: We have

$$\begin{aligned} \int_0^T \int_{\Sigma_t} |H| d\mu_t dt &= \sum_{k=0}^{T-1} \int_k^{k+1} \int_{\Sigma_t} |H| d\mu_t dt \\ &\leq \sum_{k=0}^{T-1} \left(\int_k^{k+1} \int_{\Sigma_t} |H|^2 d\mu_t dt \right)^{1/2} \left(\int_k^{k+1} \text{area} \Sigma_t \right)^{1/2} \\ &\leq \text{Area}(\Sigma_0)^{1/2} \sum_{k=0}^{T-1} \left(\int_k^{k+1} \int_{\Sigma_t} |H|^2 d\mu_t dt \right)^{1/2} \\ &\leq (C_0)^{1/2} \text{Area}(\Sigma_0)^{1/2} \sum_{k=0}^{T-1} \exp^{-\frac{Rk}{2}} \\ &\leq (C_0)^{1/2} \frac{\text{Area}(\Sigma_0)^{1/2}}{1 - \exp^{-\frac{R}{2}}}. \end{aligned}$$

This proves the proposition. Q.E.D.

Let us recall White's local regularity theorem.

Let $H(X, X_0, t)$ be the backward heat kernel on R^4 . Define

$$\rho(X, t) = 4\pi(t_0 - t)H(X, X_0, t) = \frac{1}{4\pi(t_0 - t)} \exp\left(-\frac{|X - X_0|^2}{4(t_0 - t)}\right)$$

for $t < t_0$. Let i_M be the injective radius of M^4 . We choose a cut off function $\phi \in C_0^\infty(B_{2r}(X_0))$ with $\phi \equiv 1$ in $B_r(X_0)$, where $X_0 \in M$, $0 < 2r < i_M$. Choose a normal coordinates in $B_{2r}(X_0)$ and express F using the coordinates (F^1, F^2, F^3, F^4) as a surface in R^4 . The parabolic density of the mean curvature flow is defined by

$$\Phi(X_0, t_0, t) = \int_{\Sigma_t} \phi(F) \rho(F, t) d\mu_t.$$

The following local regularity theorem was proved by White [W] (Theorem 3.1 and Theorem 4.1).

Theorem 2.3. *There is a positive constant $\varepsilon_0 > 0$ such that if*

$$\Phi(X_0, t_0, t_0 - r^2) \leq 1 + \varepsilon_0,$$

then the second fundamental form $A(t)$ of Σ_t in M is bounded in $B_{\frac{r}{2}}(X_0)$, i.e.

$$\sup_{B_{\frac{r}{2}} \times (t_0 - r^2/4, t_0]} |A| \leq C,$$

where C is a positive constant depending only on M .

Remark 2.4. *Since Σ_0 is smooth, it is well known that*

$$\lim_{r \rightarrow 0} \int_{\Sigma_0} \phi(F) \frac{1}{4\pi r^2} \exp \frac{|F - X_0|^2}{4r^2} d\mu_0 = 1$$

for any $X_0 \in \Sigma_0$. So we can find sufficiently small r_0 such that

$$\int_{\Sigma_0} \phi(F) \frac{1}{4\pi r_0^2} \exp \frac{|F - X_0|^2}{4r_0^2} d\mu_0 \leq 1 + \varepsilon_0/2$$

for all $X_0 \in M$, where ε_0 is constant in White's Theorem.

Now we state and prove our main result in this paper.

Theorem 2.5. *Suppose that M is a Kähler-Einstein surface with positive scalar curvature R . Let α be the Kähler angle of the surface Σ_t which evolves by the mean curvature flow. Then there exist sufficiently small constant ε_1 such that if $C_0 \leq \varepsilon_1$ and $\varepsilon_1/r_0^6 \ll \varepsilon_0$, where C_0 is defined in Proposition 2.1, r_0 is defined in Remark (2.4) and ε_0 is constant in White's theorem, the mean curvature flow with the initial surface Σ_0 exists globally and it converges to a holomorphic curve.*

Proof: Fix any positive T . By the definition of Φ we have

$$\Phi(X_0, t, t - r^2) = \int_{\Sigma_{t-r^2}} \phi(F) \frac{1}{4\pi r^2} \exp^{-\frac{|F - X_0|^2}{4r^2}} d\mu_{t-r^2}.$$

Differentiating this equation with respect to t we get that

$$\begin{aligned} \frac{\partial}{\partial t} \int_{\Sigma_t} \phi(F) \frac{1}{4\pi r^2} \exp^{-\frac{|F - X_0|^2}{4r^2}} d\mu_t &= \int_{\Sigma_t} \nabla \phi \cdot H \frac{1}{4\pi r^2} \exp^{-\frac{|F - X_0|^2}{4r^2}} d\mu_t \\ &\quad - \int_{\Sigma_t} \frac{\phi}{8\pi r^4} \exp^{-\frac{|F - X_0|^2}{4r^2}} \langle F - X_0, H \rangle d\mu_t \\ &\quad - \int_{\Sigma_t} \frac{\phi}{4\pi r^2} \exp^{-\frac{|F - X_0|^2}{4r^2}} |H|^2 d\mu_t. \end{aligned}$$

Integrating the above equation from r^2 to T we get that

$$\begin{aligned}
\int_{\Sigma_{T-r^2}} \phi(F) \frac{1}{4\pi r^2} \exp^{-\frac{|F-X_0|^2}{4r^2}} d\mu_t &= \int_{\Sigma_0} \phi(F) \frac{1}{4\pi r^2} \exp^{-\frac{|F-X_0|^2}{4r^2}} d\mu_0 \\
&= \int_{r^2}^T \int_{\Sigma_t} \nabla \phi \cdot H \frac{1}{4\pi r^2} \exp^{-\frac{|F-X_0|^2}{4r^2}} d\mu_t \\
&\quad - \int_{r^2}^T \int_{\Sigma_t} \frac{\phi}{8\pi r^4} \exp^{-\frac{|F-X_0|^2}{4r^2}} \langle F - X_0, H \rangle d\mu_t \\
&\quad - \int_{r^2}^T \int_{\Sigma_t} \frac{\phi}{4\pi r^2} \exp^{-\frac{|F-X_0|^2}{4r^2}} |H|^2 d\mu_t.
\end{aligned}$$

Set $r = r_0$ then we get

$$\begin{aligned}
\int_{\Sigma_{T-r_0^2}} \phi(F) \frac{1}{4\pi r_0^2} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} d\mu_t &\leq \int_{\Sigma_0} \phi(F) \frac{1}{4\pi r_0^2} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} d\mu_0 \\
&\quad + \int_0^T \int_{\Sigma_t} |\nabla \phi| |H| \frac{1}{4\pi r_0^2} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} d\mu_t \\
&\quad - \int_0^T \int_{\Sigma_t} \frac{\phi}{8\pi r_0^4} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} |F - X_0| |H| d\mu_t \\
&\quad - \int_0^T \int_{\Sigma_t} \frac{\phi}{4\pi r_0^2} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} |H|^2 d\mu_t.
\end{aligned}$$

Using Remark (2.4), Proposition 2.1, Proposition 2.2 and note that $|\nabla \phi| \leq C$, we obtain that

$$\begin{aligned}
\int_{\Sigma_{T-r_0^2}} \phi(F) \frac{1}{4\pi r_0^2} \exp^{-\frac{|F-X_0|^2}{4r_0^2}} d\mu_t &\leq 1 + \varepsilon_0/2 + \frac{C}{4\pi r_0^2} (C_0)^{1/2} \frac{Area(\Sigma_0)^{1/2}}{1 - \exp \frac{-R}{2}} \\
&\quad + \frac{1}{8\pi r_0^3} (C_0)^{1/2} \frac{Area(\Sigma_0)^{1/2}}{1 - \exp \frac{-R}{2}} + \frac{1}{4\pi r_0^2} C_0 \\
&\leq 1 + \varepsilon_0.
\end{aligned}$$

Applying White's theorem we obtain an uniform estimate of the second fundamental form which implies the global existence and convergence of the mean curvature flow. By Proposition 2.1 we can see that

$$\int_{\Sigma_t} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_t \leq C_0 \exp^{-Rt}.$$

Let $t \rightarrow \infty$ and note that $R > 0$, we get

$$\int_{\Sigma_\infty} \frac{\sin^2 \alpha}{\cos \alpha} d\mu_\infty = 0.$$

It follows that $\cos \alpha_\infty = 1$, that is Σ_∞ is a holomorphic curve. This proves the theorem. Q.E.D.

REFERENCES

- [**CL1**]: J.Chen and J.Li, Mean curvature flow of surface in 4-manifolds, *Adv. Math.* **163** (2001), 287-309.
- [**CL2**]: J. Chen and J. Li, Singularities of codimension two mean curvature flow of symplectic surfaces, preprint .
- [**CLT**]: J. Chen, J. Li and G. Tian, Two-dimensional graphs moving by mean curvature flow, *Acta Math. Sinica, English Series* Vol.**18**, No.2 (2002), 209-224.
- [**CT**]: J. Chen and G. Tian, Moving symplectic curves in Kähler-Einstein surfaces, *Acta Math. Sinica, English Series*, **16** (4), (2000), 541-548.
- [**CW**]: S.S.Chen and R.Wolfson, Minimal surfaces by moving frames, *Amer. J. Math.* **105** (1983), 59-83.
- [**EH**]: K. Ecker and G. Huisken, *Mean curvature evolution of entire graphs*, *Ann. Math.* **130** (1989), 453-471.
- [**H**]: G. Huisken, Contracting convex hypersurfaces in Riemannian manifolds by their mean curvature, *Invent. math.* **84** (1986), 463-480.
- [**Sh**]: V. Shevchishin, Pseudo-holomorphic curves and the symplectic isotopy problem, preprint.
- [**ST**]: B. Siebert and G. Tian, Holomorphy of genus two Lefschetz fibration, preprint.
- [**T**]: G. Tian, Symplectic isotopy in four dimension, First International Congress of Chinese Mathematicians (Beijing, 1998), 143-147, *AMS/IP Stud. Adv. Math.*, 20, Amer. Math. Soc., Providence, RI, 2001.
- [**W**]: B.White, A local regularity theorem for classical mean curvature flow, Preprint, Stanford.
- [**Wa**]: M.T.Wang, Mean curvature flow of surfaces in Einstein four manifolds, *J. Diff. Geom.* **57** (2001), 301-338.

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCES, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, P. R. OF CHINA.

E-mail address: hxiaoli@amss.ac.cn

MATH. GROUP, THE ABDUS SALAM ICTP, TRIESTE 34100, ITALY, AND ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCES, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, P. R. OF CHINA.

E-mail address: lijia@amss.ac.cn